

PA Description	Tranexamic Acid - Tablets (Lysteda®)
Managed by	Managed Care Organizations (MCOs) contracted by the Puerto Rico Health Insurance Administration (known in Spanish as <i>Administración de Seguros de Salud</i> or ASES) to provide pharmacy services to the insured of the Government Health Plan.
Covered Uses	All FDA approved indications: (a) The physician must document the following on the prescription: <ul style="list-style-type: none"> i. Diagnosis of hemophilia for short-term use (two to eight days) to reduce or prevent hemorrhage and reduce the need for replacement therapy during and following tooth extraction and other invasive dental procedures (ICD-10 CM: D66, D67, D68.1, D68.2, D69.1). ii. Diagnosis of treatment of cyclic heavy menstrual bleeding. (ICD-10 CM: N92.0, N92.1). iii. Heavy menstrual bleeding in females with bleeding disorders
Exclusion Criteria	(a) Using combination hormonal contraception (b) known to have any of the following conditions: <ul style="list-style-type: none"> i. Active thromboembolic disease (e.g., deep vein thrombosis, pulmonary embolism, or cerebral thrombosis) ii. A history of thrombosis or thromboembolism, including retinal vein or artery occlusion iii. An intrinsic risk of thrombosis or thromboembolism (e.g., thrombogenic valvular disease, thrombogenic cardiac rhythm disease, or hypercoagulopathy)
Required Medical Information	(a) Diagnosis of hemophilia for short-term use (two to eight days) due to <ul style="list-style-type: none"> i. Tooth extraction, OR ii. Other invasive dental procedures, OR iii. Tooth fall <p style="text-align: center;">OR</p> (b) Adjunctive treatment for mucosal bleeds (nasal, buccal or anal bleeds) (c) For diagnosis cyclic heavy menstrual bleeding: <ul style="list-style-type: none"> i. Physician certifies that patient tried and failed hormonal therapy or is contraindicated, OR ii. That patient has a diagnosed bleeding disorder.
Age Restriction	(a) None
Prescriber Restriction	(a) Hematologist (b) Hemophilia Treaters (c) Gynecologists
Coverage Duration	(a) None

Other Criteria	<p>(a) Tranexamic Acid not be administered with Factor IX Complex concentrates (For example Feiba®) or Anti-Inhibitor Coagulant concentrates, as the risk of thrombosis may be increased.</p> <p>(b) Follow Package insert instructions for dose administration.</p>
-----------------------	--